

Filtered restriction of geometric \mathcal{D} -modules

Rémi Arcadias

Depto. de Álgebra, Fac. Matemáticas, Campus Reina Mercedes
41012 Sevilla (Spain)
E-mail: rarcadias@us.es

January 23, 2012

Introduction

Let \mathcal{D} (resp. \mathcal{D}') be the ring of germs at the origin of linear partial differential operators with analytic coefficients on \mathbb{C}^n (resp. $\mathbb{C}^n \times \mathbb{C}^p$). We take a system of coordinates (x_1, \dots, x_n) on \mathbb{C}^n and $(x_1, \dots, x_n, t_1, \dots, t_p)$ on $\mathbb{C}^n \times \mathbb{C}^p$ and we denote by $i : \mathbb{C}^n \rightarrow \mathbb{C}^n \times \mathbb{C}^p$ the inclusion $x \rightarrow (x, 0)$. Let M be a finitely generated \mathcal{D}' -module specializable along $Y = \{t_1 = \dots = t_p = 0\}$. The \mathcal{D} -module theoretic restriction i^*M of M along Y is defined as the complex $(\mathcal{D}' / \sum t_i \mathcal{D}') \otimes^L M$, realized as a Koszul complex $(M \otimes \Lambda^i \mathbb{C}^p)_i$, whose cohomology groups are finitely generated over \mathcal{D} . T. Oaku and N. Takayama show in [10] how to realize this complex by a complex composed of finitely generated \mathcal{D} -modules. Let $V(\mathcal{D}')$ denote the V -filtration of Malgrange-Kashiwara along Y , and let k_1 be the maximal integral root of the b -function of M along Y . Then $(M \otimes \Lambda^i \mathbb{C}^p)_i$ is quasi-isomorphic to the sub-complex $(V_{k_1+i}(M) \otimes \Lambda^i \mathbb{C}^p)_i$. Then a free resolution of i^*M can be computed as follows: let

$$\cdots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow M \rightarrow 0 \quad (1)$$

be a free resolution of M adapted to the V -filtration. Then $(V_{k_1+i}(M) \otimes \Lambda^i \mathbb{C}^p)_i$ is isomorphic, in the derived category of \mathcal{D} -modules, to the complex

$$\cdots \rightarrow \frac{V_{k_1}(\mathcal{L}_1)}{\sum t_i V_{k_1+1}(\mathcal{L}_1)} \rightarrow \frac{V_{k_1}(\mathcal{L}_0)}{\sum t_i V_{k_1+1}(\mathcal{L}_0)} \rightarrow 0,$$

denoted by $(V_{k_1}(\mathcal{L}_i) / (\sum_j t_j V_{k_1+1}(\mathcal{L}_{i+1})))_i$.

On the other hand, the most classical filtration in \mathcal{D} -module theory is that by the order, denoted by $(F_d(\mathcal{D}))$. One defines the notion of a minimal filtered free resolution of a F -filtered \mathcal{D} -module, thus one has the notion of Betti numbers of such a module. We remark that the notion of Betti numbers can be defined as well for complexes of \mathcal{D} -modules in the derived category of F -filtered \mathcal{D} -modules $DF(\mathcal{D})$. Furthermore, T. Oaku and N. Takayama[11] and M. Granger and T. Oaku[6] define the notion of a minimal bifiltered free resolution of a

(F, V) -bifiltered \mathcal{D}' -module. With the help of this theory, we establish that the isomorphism

$$(V_{k_1+i}(M) \otimes \Lambda^i \mathbb{C}^p)_i \simeq (V_{k_1}(\mathcal{L}_i) / (\sum_j t_j V_{k_1+1}(\mathcal{L}_{i+1})))_i$$

can be seen in the category $DF(\mathcal{D})$ (Proposition 3.1).

A further property on complexes of F -filtered \mathcal{D} -modules is that of strictness, a property any filtered free resolution of a F -filtered module satisfies. In the case where $p = 1$ and $t : M \rightarrow M$ is injective, we give conditions such that the complex

$$(V_{k_1+i}(M) \otimes \Lambda^i \mathbb{C}^p)_i = (0 \rightarrow V_{k_1+1}(M) \xrightarrow{t} V_{k_1}(M) \rightarrow 0)$$

is strict (Proposition 3.3). In that case the complex

$$(V_{k_1}(\mathcal{L}_i) / (tV_{k_1+1}(\mathcal{L}_{i+1})))_i$$

becomes a F -filtered free resolution of the module M/tM . This latter fact, suggested by Toshinori Oaku, was the original motivation of this paper.

In the last section, we apply our results to the algebraic local cohomology module $N = \mathcal{O}[1/f]/\mathcal{O}$, seen as a \mathcal{D} -module. Here f is a quasi-homogeneous polynomial with an isolated singularity at the origin. N is endowed with the good F -filtration

$$F_d(N) = \sum_{1 \leq j \leq k} F_{d-j-1}(\mathcal{D})[1/f^j],$$

which takes into account the order of the pole f , where $-k$ is the least integral root of the Bernstein-Sato polynomial associated with f . By using the restriction, we give a minimal presentation associated with this data (Proposition 4.3).

1 Free resolutions of \mathcal{D} -modules

Let $\mathcal{D} = \mathbb{C}\{x_1, \dots, x_n\}[\partial_1, \dots, \partial_n]$ denote the ring of germs at the origin of linear partial differential operators with analytic coefficients on \mathbb{C}^n . It is endowed with a filtration by the order, denoted by $(F_d(\mathcal{D}))_{d \in \mathbb{Z}}$.

Let M be a finitely generated left module over \mathcal{D} . An F -filtration $(F_d(M))_{d \in \mathbb{Z}}$ on M is an exhaustive sequence of subspaces satisfying $F_d(\mathcal{D})F_{d'}(M) \subset F_{d+d'}(M)$ for any d, d' . It is called a good F -filtration if moreover there exist $f_1, \dots, f_r \in M$ and a vector shift $\mathbf{n} = (n_1, \dots, n_r)$ such that for any d ,

$$F_d(M) = \sum F_{d-n_i}(\mathcal{D})f_i.$$

For example, we denote by $\mathcal{D}^r[\mathbf{n}]$ the free module \mathcal{D}^r with basis e_1, \dots, e_r endowed with the filtration

$$F_d(\mathcal{D}^r[\mathbf{n}]) = \sum F_{d-n_i}(\mathcal{D})e_i.$$

Thus a good F -filtration of M is defined by a surjective map $\mathcal{D}^r[\mathbf{n}] \rightarrow M$.

One then defines the notion of a F -filtered free resolution: it is an exact sequence

$$\cdots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow M \rightarrow 0 \quad (2)$$

which induces for each d an exact sequence of vector spaces

$$\cdots \rightarrow F_d(\mathcal{L}_1) \rightarrow F_d(\mathcal{L}_0) \rightarrow F_d(M) \rightarrow 0.$$

Following T. Oaku and N. Takayama[11] and M. Granger and T. Oaku[6], we are able to define the notion of a *minimal* F -filtered free resolution. To that end we introduce, following F.J. Castro-Jiménez and L. Narváez-Macarro[4], the homogenization ring $\mathbf{R}\mathcal{D} = \bigoplus F_d(\mathcal{D})T^d$. It is isomorphic to the ring

$$\mathcal{D}^{(h)} = \mathbb{C}\{\mathbf{x}\}[\partial_1, \dots, \partial_n, h]$$

satisfying for any i , $\partial_i x_i - x_i \partial_i = h$. The ring $\mathcal{D}^{(h)}$ is graded by the order in ∂, h . The homogenization of a F -filtered module M is defined by

$$\mathbf{R}M = \bigoplus_d F_d(M)T^d.$$

In particular we have an isomorphism

$$\mathbf{R}(\mathcal{D}^r[\mathbf{n}]) \simeq (\mathcal{D}^{(h)})^r[\mathbf{n}]$$

where at the right hand side the vector shift \mathbf{n} refers to the grading of the module. The F -filtered free resolution (2) induces a graded free resolution

$$\cdots \rightarrow \mathbf{R}\mathcal{L}_1 \rightarrow \mathbf{R}\mathcal{L}_0 \rightarrow \mathbf{R}M \rightarrow 0. \quad (3)$$

This resolution is called *minimal* if all the entries of the matrices representing the maps $\mathbf{R}\mathcal{L}_i \rightarrow \mathbf{R}\mathcal{L}_{i-1}$ belong to the maximal two-sided ideal of $\mathcal{D}^{(h)}$ generated by $(x_i)_i, (\partial_i)_i, h$. The resolution (2) is called minimal if it induces a minimal graded free resolution (3). There exists a minimal F -filtered free resolution of M , unique up to F -filtered isomorphism (see [6]).

Let us denote now $\mathcal{D}_{x,t} = \mathbb{C}\{x_1, \dots, x_n, t_1, \dots, t_p\}[\partial_1, \dots, \partial_n, \partial_{t_1}, \dots, \partial_{t_p}]$ the ring of germs of linear partial differential operators with analytic coefficients on $\mathbb{C}^n \times \mathbb{C}^p$. It is endowed with the F -filtration by the order $(F_d(\mathcal{D}_{x,t}))$. Another filtration is the so-called V -filtration of B. Malgrange and M. Kashiwara, denoted by $(V_k(\mathcal{D}_{x,t}))_{k \in \mathbb{Z}}$, defined as follows: let us define the V -order of a monomial

$$\text{ord}^V(\lambda \mathbf{x}^\alpha t^\mu \partial^\beta \partial_t^\nu) = \sum \nu_i - \sum \mu_i,$$

with $\lambda \in \mathbb{C} \setminus 0$, then $V_k(\mathcal{D}_{x,t})$ denote the set of operators whose development only contains monomials $\mathbf{x}^\alpha t^\nu \partial^\beta \partial_t^\mu$ having order at most k .

We have also the notion of a V -filtration of a $\mathcal{D}_{x,t}$ -module: that is an exhaustive sequence of subspaces $(V_k(M))_{k \in \mathbb{Z}}$ such that for any k, k' , we have

$V_k(\mathcal{D}_{x,t})V_{k'}(M) \subset V_{k+k'}(M)$. As above let $\mathcal{D}_{x,t}^r[\mathbf{m}]$ denote the free module $\mathcal{D}_{x,t}^r$ with basis e_1, \dots, e_r endowed with the V -filtration

$$V_k(\mathcal{D}_{x,t}^r[\mathbf{m}]) = \sum V_{k-m_i}(\mathcal{D}_{x,t})e_i.$$

If M is endowed with a V -filtration defined from a surjective map $\mathcal{D}_{x,t}^r[\mathbf{m}] \rightarrow M$, then M admits a V -filtered free resolution: that is an exact sequence

$$\dots \rightarrow \mathcal{D}_{x,t}^{r_1}[\mathbf{m}^{(1)}] \rightarrow \mathcal{D}_{x,t}^{r_0}[\mathbf{m}^{(0)}] \rightarrow M \rightarrow 0$$

which induces for each k an exact sequence of vector spaces

$$\dots \rightarrow V_k(\mathcal{D}_{x,t}^{r_1}[\mathbf{m}^{(1)}]) \rightarrow V_k(\mathcal{D}_{x,t}^{r_0}[\mathbf{m}^{(0)}]) \rightarrow V_k(M) \rightarrow 0.$$

A V -filtered free resolution is the object needed in the computation of the restriction, as we shall see in the next section.

Let us describe now how to define the notion of a minimal V -filtered free resolution of M . To that end we introduce the bifiltration

$$F_{d,k}(\mathcal{D}_{x,t}) = F_d(\mathcal{D}_{x,t}) \cap V_k(\mathcal{D}_{x,t})$$

for each $d, k \in \mathbb{Z}$. A bifiltration of a $\mathcal{D}_{x,t}$ -module M is an exhaustive sequence of subspaces $(F_{d,k}(M))$ satisfying $F_{d,k}(M) \subset F_{d+1,k}(M) \cap F_{d,k+1}(M)$ and $F_{d,k}(\mathcal{D}_{x,t})F_{d',k'}(M) \subset F_{d+d',k+k'}(M)$ for any d, d', k, k' . It is called a good bifiltration if moreover there exist $f_1, \dots, f_r \in M$ and two vector shifts $\mathbf{n} = (n_1, \dots, n_r)$ and $\mathbf{m} = (m_1, \dots, m_r)$ such that for any d, k ,

$$F_{d,k}(M) = \sum F_{d-n_i, k-m_i}(\mathcal{D}_{x,t})f_i.$$

For example, we denote by $\mathcal{D}^r[\mathbf{n}][\mathbf{m}]$ the free module $\mathcal{D}_{x,t}^r$ with basis e_1, \dots, e_r endowed with the good bifiltration

$$F_{d,k}(\mathcal{D}_{x,t}^r[\mathbf{n}][\mathbf{m}]) = \sum F_{d-n_i, k-m_i}(\mathcal{D}_{x,t})e_i.$$

Let us point out that if M is endowed with a good bifiltration $(F_{d,k}(M))$, then it is endowed with a good F -filtration $F_d(M) = \cup_k F_{d,k}(M)$ and a V -filtration $V_k(M) = \cup_d F_{d,k}(M)$. A bifiltered free resolution of M is an exact sequence

$$\dots \rightarrow \mathcal{D}_{x,t}^{r_1}[\mathbf{n}^{(1)}][\mathbf{m}^{(1)}] \rightarrow \mathcal{D}_{x,t}^{r_0}[\mathbf{n}^{(0)}][\mathbf{m}^{(0)}] \rightarrow M \rightarrow 0$$

which induces for each d, k an exact sequence of vector spaces

$$\dots \rightarrow F_{d,k}(\mathcal{D}_{x,t}^{r_1}[\mathbf{n}^{(1)}][\mathbf{m}^{(1)}]) \rightarrow F_{d,k}(\mathcal{D}_{x,t}^{r_0}[\mathbf{n}^{(0)}][\mathbf{m}^{(0)}]) \rightarrow F_{d,k}(M) \rightarrow 0.$$

The ring $\mathcal{D}_{x,t}^{(h)} = \mathbf{R}\mathcal{D}_{x,t}$ is endowed by a V -filtration $V_k(\mathcal{D}_{x,t}^{(h)})$ similarly as above, by giving the weight -1 to each t_i and the weight 1 to each ∂_{t_i} . Equivalently, we have $V_k(\mathcal{D}_{x,t}^{(h)}) = \bigoplus_d V_k(\mathcal{D}_{x,t})T^d$. In the same way we endow $\mathbf{R}M$ with the

V -filtration $V_k(\mathbf{R}M) = \bigoplus_d F_{d,k}(M)T^d$. For example, we have an isomorphism of V -filtered graded $\mathcal{D}_{x,t}^{(h)}$ -modules

$$\mathbf{R}(\mathcal{D}_{x,t}^{(r)}[\mathbf{n}][\mathbf{m}]) \simeq (\mathcal{D}_{x,t}^{(h)})^r[\mathbf{n}][\mathbf{m}]$$

where the vector shift $[\mathbf{n}]$ (resp. $[\mathbf{m}]$) refers to the grading (resp. V -filtration).

Let us take a bifiltered free resolution

$$\cdots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow M \rightarrow 0. \quad (4)$$

It induces a V -filtered graded free resolution

$$\cdots \rightarrow \mathbf{R}\mathcal{L}_1 \rightarrow \mathbf{R}\mathcal{L}_0 \rightarrow \mathbf{R}M \rightarrow 0,$$

thus a bigraded free resolution

$$\cdots \rightarrow \mathrm{gr}^V(\mathbf{R}\mathcal{L}_1) \rightarrow \mathrm{gr}^V(\mathbf{R}\mathcal{L}_0) \rightarrow \mathrm{gr}^V(\mathbf{R}M) \rightarrow 0. \quad (5)$$

Since it makes sense to talk about a minimal bigraded free resolution as (4), the authors of [6] adopt the following definition: a bifiltered free resolution (4) is said to be minimal if so is the bigraded free resolution (5). They prove the existence and uniqueness (up to bifiltered isomorphism) of such a resolution.

2 Free resolutions of complexes of \mathcal{D} -modules

In that section we intend to define the notion of a minimal filtered free resolution of a complex of \mathcal{D} -modules in the corresponding derived category. We will review some facts, perhaps well-known to specialists, for the clarity of our text.

Let us denote by $CF(\mathcal{D})$ the category of bounded complexes $\cdots \rightarrow M_i \rightarrow M_{i-1} \rightarrow \cdots$ where for each i , M_i is a F -filtered \mathcal{D} -module and the differentials are F -adapted.

A map $\alpha : M_\bullet \rightarrow N_\bullet$ is said to be a filtered quasi-isomorphism if the induced map $\mathrm{gr}(\alpha) : \mathrm{gr}^F M_\bullet \rightarrow \mathrm{gr}^F N_\bullet$ is a quasi-isomorphism.

If $\alpha : M_\bullet \rightarrow N_\bullet$ is a morphism in $CF(\mathcal{D})$, we denote by $C(\alpha)$ the mapping cone of α , defined by $C(\alpha)_n = M_{n-1} \oplus N_n$ with differential $(x, y) \mapsto (-\phi(x), \psi(y) + \alpha(x))$. It is endowed with the filtration $F_d(C(\alpha)_i) = F_d(M_{i-1}) \oplus F_d(N_i)$. In that way we have natural isomorphisms $F_d(C(\alpha)) \simeq C(F_d(\alpha))$ and $\mathrm{gr}^F(C(\alpha)) \simeq C(\mathrm{gr}^F(\alpha))$.

Lemma 2.1. *Let $\alpha : M_\bullet \rightarrow N_\bullet$ as above. If for any d , $F_d(\alpha)$ is a quasi-isomorphism, then $\mathrm{gr}^F(\alpha)$ is a quasi-isomorphism.*

Proof. $F_d(\alpha)$ is a quasi-isomorphism is equivalent to saying that $C(F_d(\alpha))$ is exact. Thus $F_d(C(\alpha))$ is exact and it easily implies that $\mathrm{gr}^F(C(\alpha)) \simeq C(\mathrm{gr}^F(\alpha))$ is exact. That gives the result. \square

We have a converse statement:

Proposition 2.1. *Let $\alpha : M_\bullet \rightarrow N_\bullet$ be a morphism in $CF(\mathcal{D})$, such that for each i , M_i and N_i are endowed with good filtrations. If $\text{gr}^F(\alpha)$ is a quasi-isomorphism, then α is a quasi-isomorphism and induces for each d a quasi-isomorphism $F_d(\alpha)$.*

Proof. We have that $\text{gr}^F(C(\alpha)) \simeq C(\text{gr}^F(\alpha))$ is exact. Since each $C(\alpha)_i$ is endowed with a good filtration, then by a statement analogous to [6], Proposition 2.5, it follows that $C(\alpha)$ is exact, moreover for each d , $F_d(C(\alpha))$ is exact, which establishes the statement. \square

Corollary 2.1. *Let $\alpha : M_\bullet \rightarrow N_\bullet$ be a morphism in $CF(\mathcal{D})$, such that for each i , M_i and N_i are endowed with good filtrations. Then α is a filtered quasi-isomorphism if and only if for each d , $F_d(\alpha)$ is a quasi-isomorphism. Moreover, if α is a filtered quasi-isomorphism, then it is a quasi-isomorphism.*

A complex

$$\cdots \rightarrow M_i \xrightarrow{\phi_i} M_{i-1} \rightarrow \cdots$$

in $CF(\mathcal{D})$ is said to be *strict* if for each i and d we have $\text{Im} \phi_i \cap F_d(M_{i-1}) = \phi_i(F_d(M_i))$.

Lemma 2.2. *Let*

$$\begin{array}{ccccc} M_2 & \xrightarrow{\phi_2} & M_1 & \xrightarrow{\phi_1} & M_0 \\ \downarrow & & \downarrow \alpha_1 & & \downarrow \\ N_2 & \xrightarrow{\psi_2} & N_1 & \xrightarrow{\psi_1} & N_0 \end{array}$$

be a commutative diagram of filtered \mathcal{D} -modules, such that the rows are complexes and α_1 induces isomorphisms $H_1(M_\bullet) \simeq H_1(N_\bullet)$ and $H_1(F_d(M_\bullet)) \simeq H_1(F_d(N_\bullet))$. Then α_1 induces an isomorphism

$$\frac{F_d(\phi_2(M_2))}{\phi_2(F_d(M_2))} \simeq \frac{F_d(\psi_2(N_2))}{\psi_2(F_d(N_2))}.$$

Proof. We have a commutative diagram with exact rows

$$\begin{array}{ccccccc} \frac{F_d(\text{Ker } \phi_1)}{\phi_2(F_d(M_2))} & \longrightarrow & \frac{\text{Ker } \phi_1}{\text{Im } \phi_2} & \longrightarrow & \frac{\text{Ker } \phi_1}{F_d(\text{Ker } \phi_1) + \text{Im } \phi_2} & \longrightarrow & 0 \\ \downarrow \text{iso} & & \downarrow \text{iso} & & \downarrow & & \\ \frac{F_d(\text{Ker } \psi_1)}{\psi_2(F_d(N_2))} & \longrightarrow & \frac{\text{Ker } \psi_1}{\text{Im } \psi_2} & \longrightarrow & \frac{\text{Ker } \psi_1}{F_d(\text{Ker } \psi_1) + \text{Im } \psi_2} & \longrightarrow & 0, \end{array}$$

then by the five lemma, α_1 induces an isomorphism

$$\frac{\text{Ker } \phi_1}{F_d(\text{Ker } \phi_1) + \text{Im } \phi_2} \simeq \frac{\text{Ker } \psi_1}{F_d(\text{Ker } \psi_1) + \text{Im } \psi_2}.$$

Similarly, from the diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & \frac{F_d(\text{Ker } \phi_1)}{F_d(\text{Im } \phi_2)} & \longrightarrow & \frac{\text{Ker } \phi_1}{\text{Im } \phi_2} & \longrightarrow & \frac{\text{Ker } \phi_1}{F_d(\text{Ker } \phi_1) + \text{Im } \phi_2} \longrightarrow 0 \\
& & \downarrow & & \downarrow \text{iso} & & \downarrow \text{iso} \\
0 & \longrightarrow & \frac{F_d(\text{Ker } \psi_1)}{F_d(\text{Im } \psi_2)} & \longrightarrow & \frac{\text{Ker } \psi_1}{\text{Im } \psi_2} & \longrightarrow & \frac{\text{Ker } \psi_1}{F_d(\text{Ker } \psi_1) + \text{Im } \psi_2} \longrightarrow 0,
\end{array}$$

we deduce that α_1 induces an isomorphism

$$\frac{F_d(\text{Ker } \phi_1)}{F_d(\text{Im } \phi_2)} \simeq \frac{F_d(\text{Ker } \psi_1)}{F_d(\text{Im } \psi_2)}.$$

Similarly, from the diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & \frac{F_d(\phi_2(M_2))}{\phi_2(F_d(M_2))} & \longrightarrow & \frac{F_d(\text{Ker } \phi_1)}{\phi_2(F_d(M_2))} & \longrightarrow & \frac{F_d(\text{Ker } \phi_1)}{F_d(\phi_2(M_2))} \longrightarrow 0 \\
& & \downarrow & & \downarrow \text{iso} & & \downarrow \text{iso} \\
0 & \longrightarrow & \frac{F_d(\psi_2(N_2))}{\psi_2(F_d(N_2))} & \longrightarrow & \frac{F_d(\text{Ker } \psi_1)}{\psi_2(F_d(N_2))} & \longrightarrow & \frac{F_d(\text{Ker } \psi_1)}{F_d(\psi_2(N_2))} \longrightarrow 0,
\end{array}$$

the result follows. \square

Corollary 2.2. *Let $\alpha : M_\bullet \rightarrow N_\bullet$ be a morphism in $CF(\mathcal{D})$ which is a quasi-isomorphism and which induces quasi-isomorphisms $F_d(M_\bullet) \rightarrow F_d(N_\bullet)$. Then C_\bullet is strict if and only if D_\bullet is strict.*

Let $KF(\mathcal{D})$ denote the category whose objects are the objects of $CF(\mathcal{D})$ and the maps are taken modulo (F -adapted) homotopies. Then $DF(\mathcal{D})$ denotes the localisation of $KF(\mathcal{D})$ with respect to filtered quasi-isomorphisms, as done in [12], 2.1.8 (see also [8]). In other terms, $DF(\mathcal{D})$ is the localization of $KF(\mathcal{D})$ with respect to the null system composed of the complexes M_\bullet such that $\text{gr}(M_\bullet)$ is acyclic (see in [7] an introduction to localization of categories). $D(\mathcal{D})$ will denote the derived category of \mathcal{D} -modules.

Because of Corollary 2.2, the strictness of a complex M_\bullet , where each M_i is endowed with a good filtration, makes sense in $DF(\mathcal{D})$.

As pointed out in [13], for a strict complex M_\bullet , for any i, d , $H_i(F_d(M_\bullet))$ is a subspace of $H_i(\mathcal{M}_\bullet)$, more precisely we can define a good filtration on $H_i(\mathcal{M}_\bullet)$ by $F_d(H_i(\mathcal{M})) = H_i(F_d(M_\bullet))$.

Let \mathfrak{m} denotes the maximal graded ideal of $\text{gr}^F(\mathcal{D})$ generated by

$$x_1, \dots, x_n, \xi_1, \dots, \xi_n$$

and $\mathfrak{m}^{(h)}$ the maximal graded two-sided ideal of $\mathcal{D}^{(h)}$ generated by

$$x_1, \dots, x_n, \partial_1, \dots, \partial_n, h.$$

We have $\text{gr}^F(\mathcal{D})/\mathfrak{m} \simeq \mathbb{C}$ and $\mathcal{D}^{(h)}/\mathfrak{m}^{(h)} \simeq \mathbb{C}$.

Definition 2.1. Let $\mathcal{L}_\bullet = (\cdots \rightarrow \mathcal{L}_i \xrightarrow{\psi_i} \mathcal{L}_{i-1} \rightarrow \cdots)$ a complex of $CF(\mathcal{D})$ with for each i , $\mathcal{L}_i = \mathcal{D}^{r_i}[\mathbf{n}^{(i)}]$. The complex \mathcal{L}_\bullet is said to be minimal if for any i , the matrix representing $\mathbf{R}\psi_i$ as its entries in $\mathfrak{m}^{(h)}$ (equivalently, for any i , the matrix representing $\text{gr}\psi_i$ as its entries in \mathfrak{m}).

Definition 2.2. Let M_\bullet be a complex of $DF(\mathcal{D})$. A minimal resolution of M_\bullet is a minimal complex \mathcal{L}_\bullet isomorphic to M_\bullet in $DF(\mathcal{D})$.

The numbers $r^{(i)}$ and the shifts $\mathbf{n}^{(i)}$ arising in a minimal resolution of M_\bullet make sense. In fact, denoting

$$\begin{aligned}\beta_{i,j} &= \dim_{\mathbb{C}} \text{Tor}_i^{\text{gr}^F(\mathcal{D})}(\text{gr}^F(M), \mathbb{C})_j \\ &= \dim_{\mathbb{C}} \text{Tor}_i^{\mathcal{D}^{(h)}}(\mathcal{R}_F(M), \mathbb{C})_j,\end{aligned}$$

we have $\beta_{i,j} = \text{card}\{k, n_k^{(i)} = j\}$. In particular $r_i = \sum_j \beta_{i,j}$.

Let us see finally the link between resolutions of modules and resolutions of complexes. The notion of a strict filtered free resolution of a complex generalizes the notion of a filtered free resolution of a module: regarding a filtered module M as a complex in $DF(\mathcal{D})$ concentrated in degree 0, a strict filtered free resolution $\cdots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow 0$ of M (thus $H_0(\mathcal{L}_\bullet) \simeq M$) provides a filtered free resolution $\cdots \rightarrow \mathcal{L}_1 \rightarrow \mathcal{L}_0 \rightarrow H_0(\mathcal{L}_\bullet) \rightarrow 0$ in the sense of [11].

3 Filtered restriction

Let M be a $\mathcal{D}_{x,t}$ -module endowed with a good bibiltration $(F_{d,k}(M))$ such that $M = \mathcal{D}_{x,t}V_0(M)$. We assume that there exists a non-zero polynomial $b(s) \in \mathbb{C}[s]$ such that $b(t_1\partial_{t_1} + \cdots + t_p\partial_{t_p})\text{gr}_0^V(M) = 0$, which is the case if M is holonomic. An algorithm to compute such a polynomial can be found in [10]. Let k_1 be an integer such that $b(k) \neq 0$ if $k > k_1$.

Let $i : (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}^n \times \mathbb{C}^p, 0)$ denote the embedding $x \rightarrow (x, 0) = (x, t)$. The restriction i^*M of M along $Y = \{t_1 = \cdots = t_p = 0\}$ is by definition the complex $(\mathcal{D}_{x,t}/\sum_i T_i \mathcal{D}_{x,t}) \otimes_{\mathcal{D}_{x,t}}^L M$ in $D(\mathcal{D})$ -modules. Let $\Lambda^i = \Lambda^i \mathbb{C}^p$. Then i^*M is represented by the Koszul complex over the sequence t_1, \dots, t_p :

$$0 \rightarrow M \otimes \Lambda^p \xrightarrow{\delta} \cdots \rightarrow M \otimes \Lambda^1 \xrightarrow{\delta} M \otimes \Lambda^0 \rightarrow 0.$$

This complex is made of non finitely generated \mathcal{D} -modules. By [9], Proposition 5.2 and [10], section 5, the truncation

$$0 \rightarrow V_{k_1+p}(M) \otimes \Lambda^p \xrightarrow{\delta} \cdots \rightarrow V_{k_1+1}(M) \otimes \Lambda^1 \xrightarrow{\delta} V_{k_1}(M) \otimes \Lambda^0 \rightarrow 0,$$

denoted by $V_{k_1+\bullet} \otimes \Lambda^\bullet$, is quasi-isomorphic to the above Koszul complex, thus still represents i^*M . The bifiltration allows us to endow $V_{k_1+\bullet} \otimes \Lambda^\bullet$ with a filtration $F_d(V_{k_1+\bullet} \otimes \Lambda^\bullet)$:

$$0 \rightarrow F_{d,k_1+p}(M) \otimes \Lambda^p \xrightarrow{\delta} \cdots \rightarrow F_{d,k_1+1}(M) \otimes \Lambda^1 \xrightarrow{\delta} F_{d,k_1}(M) \otimes \Lambda^0 \rightarrow 0.$$

Let us remark that since the bifiltration of M is good, then there exists f_1, \dots, f_r such that $F_{d,k}(M) = \sum_i F_{d-n_i, k-m_i}(\mathcal{D})f_i$. If k_1 is chosen so that for any i , $k_1 \geq m_i$, then each f_i belongs to $V_{k_1}(M)$ and for each $i = 0, \dots, p$, $F_{d, k_1+i}(M)$ is a good filtration of $V_{k_1+i}(M)$.

Let

$$\cdots \rightarrow \mathcal{L}_1 \xrightarrow{\psi_1} \mathcal{L}_0 \xrightarrow{\psi_0} M \rightarrow 0$$

be any bifiltered free resolution of M , with $\mathcal{L}_i = \mathcal{D}_{x,t}^{r(i)}[\mathbf{n}^{(i)}][\mathbf{m}^{(i)}]$. In particular it is a V -filtered free resolution. By [10], Theorem 5.3, the complex

$$\cdots \rightarrow \frac{V_{k_1}(\mathcal{L}_1)}{\sum_i t_i V_{k_1+1}(\mathcal{L}_1)} \rightarrow \frac{V_{k_1}(\mathcal{L}_0)}{\sum_i t_i V_{k_1+1}(\mathcal{L}_0)} \rightarrow 0,$$

denoted by $V_{k_1}(\mathcal{L}_\bullet)/\sum_i t_i V_{k_1+1}(\mathcal{L}_\bullet)$ is a free complex isomorphic to i^*M in $D(\mathcal{D})$. We endow it with a F -filtration by setting

$$\begin{aligned} F_d \left(\frac{V_{k_1}(\mathcal{L}_i)}{tV_{k_1+1}(\mathcal{L}_i)} \right) &= \frac{F_{d,k_1}(\mathcal{L}_i)}{F_{d,k_1}(\mathcal{L}_i) \cap tV_{k_1+1}(\mathcal{L}_i)} \\ &\simeq \frac{F_{d,k_1}(\mathcal{L}_i)}{tF_{d,k_1+1}(\mathcal{L}_i)}. \end{aligned}$$

Note that the filtered \mathcal{D} -module $V_{k_1}(\mathcal{L}_i)/\sum_i t_i V_{k_1+1}(\mathcal{L}_i)$ is naturally isomorphic to some $\mathcal{D}^r[\mathbf{n}]$.

Proposition 3.1. *The complexes $V_{k_1+\bullet}(M) \otimes \Lambda^\bullet$ and $V_{k_1}(\mathcal{L}_\bullet)/\sum_i t_i V_{k_1+1}(\mathcal{L}_\bullet)$ are isomorphic in $DF(\mathcal{D})$. Thus $V_{k_1}(\mathcal{L}_\bullet)/\sum_i t_i V_{k_1+1}(\mathcal{L}_\bullet)$ is a filtered free resolution of i^*M .*

Proof. Since t_1, \dots, t_p is a regular sequence in $\mathcal{O}_{x,t}$, we have that the Koszul complex

$$0 \rightarrow \mathbf{R}_{F,V}(\mathcal{D}_{x,t}) \otimes \Lambda^p \rightarrow \cdots \rightarrow \mathbf{R}_{F,V}(\mathcal{D}_{x,t}) \otimes \Lambda^0 \rightarrow \frac{\mathbf{R}_{F,V}(\mathcal{D}_{x,t})}{\sum_i t_i \mathbf{R}_{F,V}(\mathcal{D}_{x,t})} \rightarrow 0$$

is exact. We may replace $\mathcal{D}_{x,t}$ by the bifiltered free module \mathcal{L}_i , then the complex

$$0 \rightarrow \mathbf{R}_{F,V}(\mathcal{L}_i) \otimes \Lambda^p \xrightarrow{\delta} \cdots \rightarrow \mathbf{R}_{F,V}(\mathcal{L}_i) \otimes \Lambda^0 \rightarrow \frac{\mathbf{R}_{F,V}(\mathcal{L}_i)}{\sum_i t_i \mathbf{R}_{F,V}(\mathcal{L}_i)} \rightarrow 0$$

is exact. Moreover it is bigraded, thus the complex

$$0 \rightarrow F_{d,k_1+p}(\mathcal{L}_i) \otimes \Lambda^p \xrightarrow{\delta} \cdots \rightarrow F_{d,k_1}(\mathcal{L}_i) \otimes \Lambda^0 \rightarrow \frac{F_{d,k_1}(\mathcal{L}_i)}{\sum_i t_i F_{d,k_1+1}(\mathcal{L}_i)} \rightarrow 0$$

is exact. We have a commutative diagram:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \uparrow & & \uparrow & & \\
\cdots & \longrightarrow & F_{d,k_1+1}(M) \otimes \Lambda^1 & \longrightarrow & F_{d,k_1}(M) \otimes \Lambda^0 & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
\cdots & \longrightarrow & F_{d,k_1+1}(\mathcal{L}_0) \otimes \Lambda^1 & \longrightarrow & F_{d,k_1}(\mathcal{L}_0) \otimes \Lambda^0 & \longrightarrow & \frac{F_{d,k_1}(\mathcal{L}_0)}{\sum_i t_i F_{d,k_1+1}(\mathcal{L}_0)} \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
\cdots & \longrightarrow & F_{d,k_1+1}(\mathcal{L}_1) \otimes \Lambda^1 & \longrightarrow & F_{d,k_1}(\mathcal{L}_1) \otimes \Lambda^0 & \longrightarrow & \frac{F_{d,k_1}(\mathcal{L}_1)}{\sum_i t_i F_{d,k_1+1}(\mathcal{L}_1)} \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
& & \vdots & & \vdots & & \vdots
\end{array}$$

We deduce that the complexes $(F_{d,k_1+i}(M) \otimes \Lambda^i)_i$ and $(V_{k_1}(\mathcal{L}_i) / \sum_i t_i V_{k_1+1}(\mathcal{L}_i))_i$ are both isomorphic to the complex associated with the double complex

$$(F_{d,k_1+j}(\mathcal{L}_i) \otimes \Lambda^j)_{i,j}.$$

□

Corollary 3.1. *The complex $V_{k_1}(\mathcal{L}_\bullet) / \sum_i t_i V_{k_1+1}(\mathcal{L}_\bullet)$ is strict if so is the complex $V_{k_1+\bullet}(M) \otimes \Lambda^\bullet$.*

Let us consider from now on a special case: $p = 1$. The complex $V_{k_1+\bullet}(M) \otimes \Lambda^\bullet$ is $V_{k_1+1}(M) \xrightarrow{t} V_{k_1}(M)$. Let us assume furthermore that $t : M \rightarrow M$ is injective. The restriction i^*M is concentrated in degree 0 with $H_0 i^*M \simeq M/tM$. Thus the complex

$$\cdots \rightarrow \frac{V_{k_1}(\mathcal{L}_1)}{tV_{k_1+1}(\mathcal{L}_1)} \rightarrow \frac{V_{k_1}(\mathcal{L}_0)}{tV_{k_1+1}(\mathcal{L}_0)} \rightarrow \frac{M}{tM} \rightarrow 0 \quad (6)$$

is a free resolution of the module $M/tM = (V_{k_1}(M) + tM)/tM$. That module is naturally endowed with the filtration

$$F_d(M/tM) := \frac{F_{d,k_1}(M)}{F_{d,k_1}(M) \cap tM}.$$

and from Proposition 3.1 and Corollary 3.1 we obtain:

Proposition 3.2. *Assume that $t : M \rightarrow M$ is injective and that for any d , $F_{d,k_1}(M) \cap tV_{k_1+1}(M) = tF_{d,k_1+1}(M)$. Then the complex (6) is an F -filtered free resolution of M/tM , i.e. for any d the complex*

$$\cdots \rightarrow \frac{F_{d,k_1}(\mathcal{L}_1)}{tF_{d,k_1+1}(\mathcal{L}_1)} \rightarrow \frac{F_{d,k_1}(\mathcal{L}_0)}{tF_{d,k_1+1}(\mathcal{L}_0)} \rightarrow F_d\left(\frac{M}{tM}\right) \rightarrow 0 \quad (7)$$

is exact. The homogenization, with respect to F , of the resolution (6) is as follows:

$$\cdots \rightarrow \frac{V_{k_1}(\mathbf{R}\mathcal{L}_1)}{tV_{k_1+1}(\mathbf{R}\mathcal{L}_1)} \rightarrow \frac{V_{k_1}(\mathbf{R}\mathcal{L}_0)}{tV_{k_1+1}(\mathbf{R}\mathcal{L}_0)} \rightarrow \mathbf{R}\left(\frac{M}{tM}\right) \rightarrow 0 \quad (8)$$

Let us remark that the minimality of the bifiltered free resolution $\cdots \rightarrow \mathcal{L}_0 \rightarrow M \rightarrow 0$ does not imply the minimality of the F -filtered free resolution (6), as we shall see in Section 4.

Let us give sufficient conditions for the assumptions in 3.2.

Proposition 3.3. *The assumptions in Proposition 3.2 hold if the map $t : \text{gr}^F(M) \rightarrow \text{gr}^F(M)$ is injective and for any d, k , $F_{d,k}(M) = F_d(M) \cap V_k(M)$.*

That follows from the two following lemmas.

Lemma 3.1. *Let us assume*

1. *The map $t : M \rightarrow M$ is injective,*
2. *For any d , $F_d(M) \cap tM = tF_d(M)$,*
3. *For any d, k , $F_{d,k}(M) = F_d(M) \cap V_k(M)$.*

Then for any d , $tM \cap F_{d,k_1}(M) = tF_{d,k_1+1}(M)$,

Proof. By the injectivity of equation (5.2) in [9], we have $tM \cap V_{k_1}(M) = tV_{k_1+1}(M)$. Then

$$\begin{aligned} tM \cap F_{d,k_1}(M) &\subset tV_{k_1+1}(M) \cap F_d(M) \\ &\subset tV_{k_1+1}(M) \cap tF_d(M) \quad \text{by assumption 2} \\ &\subset t(V_{k_1+1}(M) \cap F_d(M)) \quad \text{by assumption 1} \\ &\subset tF_{d,k_1+1}(M) \quad \text{by assumption 3.} \end{aligned}$$

□

Lemma 3.2. *The conditions 1. and 2. of the preceding Lemma are satisfied if and only if the map $t : \text{gr}^F(M) \rightarrow \text{gr}^F(M)$ is injective.*

Let us give more information on the assumptions of Proposition 3.3.

Lemma 3.3 ([1], Lemma 1.1). *$\forall d, k$, $F_{d,k}(M) = F_d(M) \cap V_k(M)$ holds if and only if the map induced by h on $\text{gr}^V(\mathbf{R}M)$ is injective.*

Remark 3.1. The conditions in Proposition 3.3 depend on the good bifiltration. The first condition is that $t : \text{gr}^F(M) \rightarrow \text{gr}^F(M)$ is injective. Let $M = \mathcal{D}_{x,t}^2 / (\mathcal{D}_{x,t} \cdot (t, 1))$. If we define

$$F_d(M) = F_{d-1}(\mathcal{D}_{x,t}) \cdot \overline{(1, 0)} + F_d(\mathcal{D}_{x,t}) \cdot \overline{(0, 1)},$$

then t is not injective on $\text{gr}^F(M) \simeq \text{gr}^F(\mathcal{D}_{x,t})^2/(t, 0)$. On the other hand, defining

$$F_d(M) = F_d(\mathcal{D}_{x,t}).\overline{(1, 0)} + F_{d-1}(\mathcal{D}_{x,t}).\overline{(0, 1)},$$

then t is injective on $\text{gr}^F(M) \simeq \text{gr}^F(\mathcal{D}_{x,t})^2/(0, 1)$.

The second condition is that for any d, k , $F_{d,k}(M) = F_d(M) \cap V_k(M)$, which is equivalent to the injectivity of the map induced by h on $\text{gr}^V(\mathbf{R}(M))$. Let us take $M = \mathcal{D}_{x,t}$ endowed with the usual bifiltration, then the above condition is satisfied. Let $M' = \mathcal{D}_{x,t}^2/(1, 1)$. We have an isomorphism $M \simeq M'$ given by $1 \mapsto \overline{(1, 0)}$. Let us endow M' with the good bifiltration defined by

$$F_{d,k}(M') = F_{d-1,k}(\mathcal{D}_{x,t}).\overline{(1, 0)} + F_{d,k-1}(\mathcal{D}_{x,t}).\overline{(0, 1)}.$$

Then $\mathbf{R}(M) \simeq (\mathcal{D}_{x,t}^{(h)})^2[0, 1]/(1, h)$ and $\text{gr}^V(\mathbf{R}(M)) \simeq (\text{gr}^V(\mathcal{D}_{x,t}^{(h)}))^2/(0, h)$, on which the map induced by h is not injective.

To end this section, let us recall the notion of involutive bases, which we will use in Section 4.

Definition 3.1. Let M be a filtered \mathcal{D} -module. If $0 \neq m \in M$, we define the F -order of m by $\text{ord}^F(m) = \min\{d, m \in F_d(M)\}$ and the F -symbol $\sigma^F(m)$ by the class of m in $\text{gr}_{\text{ord}^F(m)}^F(M)$. If it is clear in the context we will simply note $\sigma(m)$.

Let I be an ideal of \mathcal{D} , endowed with the induced filtration. We say that (P_1, \dots, P_r) is an F -involutive base of I if for any $P \in I$, there exist Q_1, \dots, Q_r such that $P = \sum Q_i P_i$ and for any i , $\text{ord}^F(Q_i P_i) \leq \text{ord}^F(P)$.

We have the following useful characterization of involutive bases.

Proposition 3.4. Let I be generated by P_1, \dots, P_r and let $n_i = \text{ord}^F(P_i)$. Assume that we have homogeneous elements

$$S_i = \sum_j S_{i,j} e_j \in \text{gr}^F(\mathcal{D})^r[\mathbf{n}]$$

which generate the relations between $\sigma(P_1), \dots, \sigma(P_r)$, such that for any i , there exists a relation

$$R_i = \sum_j R_{i,j} e_j \in \mathcal{D}[\mathbf{n}]$$

between P_1, \dots, P_r , such that $\sigma(R_i) = S_i$. Then (P_1, \dots, P_r) is an F -involutive base of I .

Proof. Let $P = \sum Q_j P_j$ and assume that $\text{ord}(P) < \text{ord}(\sum Q_j e_j) = d$ with d minimal. Then $\sigma(\sum Q_j P_j) = 0$, which leads to a relation $\sigma(\sum Q_j e_j)$ between the $(\sigma(P_j))$. Thus for any i there exists $a_i = \sigma(b_i)$ such that $\sigma(\sum Q_j e_j) = \sum a_i S_i = \sigma(\sum b_i R_i)$. Then

$$P = (\sum Q_j e_j - \sum b_i R_i).(P_1, \dots, P_r)$$

with $\text{ord}(\sum Q_j e_j - \sum b_i R_i) < d$, a contradiction. \square

We will say that the relation S_i is *lifted* by the relation R_i such that $\sigma(R_i) = S_i$. In fact it is easy to see that the converse of the lemma holds: if P_1, \dots, P_r is an F -involutive base, then every homogeneous relation between $(\sigma(P_i))$ can be lifted by a relation between the (P_i) .

4 Application to algebraic local cohomology

Our aim is to give presentations of some algebraic local cohomology module, viewed as a \mathcal{D} -module. Let $\mathcal{O} = \mathbb{C}\{\mathbf{x}\}$ and let $f \in \mathcal{O}$. We focus on the module $N = \mathcal{O}[1/f]/\mathcal{O}$ which is quasi-isomorphic to the complex of algebraic local cohomology $\mathbb{R}^{\bullet+1}\Gamma_{[f=0]}(\mathcal{O})$. Although this module is not finitely generated over \mathcal{O} , it turns to be a finitely generated \mathcal{D} -module. We will assume that f is quasi-homogeneous and has an isolated singularity at the origin. We will give two minimal presentations of N , one of which being classical, the other coming from our general result in Section 3.

Let $f \neq 0$ be any function in \mathcal{O} . The vector space $\mathcal{O}[1/f, s]f^s$, where f^s is understood as a symbol, has a natural structure of a $\mathcal{D}[s]$ -module. Let us recall that there exists a polynomial $b_f(s) \in \mathbb{C}[s]$ called *Bernstein-Sato polynomial* such that there exists $P(s) \in \mathcal{D}[s]$ such that

$$b(s)f^s = P(s)f^{s+1}. \quad (9)$$

Let $-k'$ be the least integral root of $b(s)$. It is known that $k' \geq 1$. Using the functional equation (9), one has $\mathcal{O}[1/f] = \mathcal{D}(1/f^{k'})$. Let us assume from now on that f has an isolated singularity at the origin. Let

$$S_{i,j} = f'_i \partial_j - f'_j \partial_i$$

for any $0 < i < j \leq n$. Then the symbols $(\sigma^F(S_{i,j}))$ generate the kernel of the map of \mathcal{O} -algebras

$$\phi_f : \mathcal{O}[\xi] \rightarrow \bigoplus (J(f))^d T^d$$

defined by $\xi_i \mapsto f'_i T$, and the operators $(S_{i,j})$ generate $\text{Ann}_{\mathcal{D}} f^s$, see [3]. Let us identify $\text{gr}^F(\mathcal{D})$ with $\mathbb{C}\{x_1, \dots, x_n\}[\xi_1, \dots, \xi_n]$. It is also known that the relations between the symbols $(\sigma(S_{i,j}))$ are generated by

$$f'_i \sigma(S_{j,k}) - f'_j \sigma(S_{i,k}) + f'_k \sigma(S_{i,j}) = 0 \quad (10)$$

and

$$\xi_i \sigma(S_{j,k}) - \xi_j \sigma(S_{i,k}) + \xi_k \sigma(S_{i,j}) = 0 \quad (11)$$

for each triple $i < j < k$ (see [1], paragraph 3.1.1). These relations are lifted by the following relations in \mathcal{D} :

$$f'_i S_{j,k} - f'_j S_{i,k} + f'_k S_{i,j} = 0$$

and

$$\partial_i S_{j,k} - \partial_j S_{i,k} + \partial_k S_{i,j} = 0.$$

We also assume that f is quasi-homogeneous, i.e. there exist positive weights w_1, \dots, w_n such that, denoting $\theta = \sum w_i x_i \partial_i$, we have $\theta(f) = f$.

4.1 Classical presentation of $\mathcal{O}[1/f]/\mathcal{O}$

First let us consider the module $\mathcal{O}[1/f]$ endowed with the following good F -filtration:

$$F_d \left(\mathcal{O} \left[\frac{1}{f} \right] \right) = F_d(\mathcal{D}) \frac{1}{f^{k'}}.$$

Proposition 4.1. *The module $\mathcal{O}[1/f]$ admits the minimal filtered presentation $\mathcal{L}_1 \xrightarrow{\phi_1} \mathcal{D}[0] \xrightarrow{\phi_0} \mathcal{O}[1/f] \rightarrow 0$ where*

- $\phi_0(1) = 1/f^{k'}$,
- $\mathcal{L}_1 = \mathcal{D}e \oplus (\bigoplus_{i < j} \mathcal{D}e_{i,j})$ with $\text{ord}(e) = 1$ and $\text{ord}(e_{i,j}) = 1$,
- $\phi_1(e) = \theta + k'$,
- $\phi_1(e_{i,j}) = S_{i,j}$.

Proof. Since $\text{Ann}_{\mathcal{D}} f^s$ is generated by $(S_{i,j})$, we have that $\text{Ann}_{\mathcal{D}[s]} f^s$ is generated by $(S_{i,j})$ and $s - \theta$. Using [5], Remark 14, we deduce that $\text{Ann}_{\mathcal{D}}(1/f^{k'})$ is generated by $(S_{i,j})$ and $\theta + k$. Let us show that this system of generators is F -involutive.

We claim that $\sigma(\theta)$ is a non-zero divisor on $\mathcal{O}[\xi]/(\sigma(S_{i,j}))$. Indeed, assume that $P\sigma(\theta) \in (S_{i,j})$. Then $0 = \phi_f(P\sigma(\theta)) = \phi_f(P)f$ thus $\phi_f(P) = 0$ and $P \in (\sigma(S_{i,j}))$ as claimed. Next, the relation

$$\sigma(\theta)\sigma(S_{i,j}) - \sigma(S_{i,j})\sigma(\theta) = 0 \tag{12}$$

can be lifted since $[\theta + k', S_{i,j}] = (1 - w_i - w_j)S_{i,j}$ which reads

$$(\theta + k' - 1 + w_i + w_j)S_{i,j} - S_{i,j}(\theta + k') = 0.$$

The involutivity follows then from Proposition 3.4, and the minimality of the presentation is clear. \square

Now we turn to the module N , which we endow similarly with the following good F -filtration:

$$F_d(N) = F_d(\mathcal{D}) \left[\frac{1}{f^{k'}} \right].$$

Proposition 4.2. *The module $N = \mathcal{O}[1/f]/\mathcal{O}$ admits the minimal filtered presentation $\mathcal{L}_1 \xrightarrow{\phi_1} \mathcal{D}[0] \xrightarrow{\phi_0} N \rightarrow 0$ where*

- $\phi_0(1) = [1/f^{k'}]$,
- $\mathcal{L}_1 = \mathcal{D}e_0 \oplus \mathcal{D}e_1 \oplus (\bigoplus_{i < j} \mathcal{D}e_{i,j})$ with $\text{ord}(e_0) = 0$, $\text{ord}(e_1) = 1$, $\text{ord}(e_{i,j}) = 1$,
- $\phi_1(e_0) = f^{k'}$,
- $\phi_1(e_1) = \theta + k'$,

- $\phi_1(e_{i,j}) = S_{i,j}$.

Proof. First, let us show that $\text{Ker}\phi_0 = \text{Im}\phi_1$. For $P \in \mathcal{D}$, $\phi_0(P) = 0$ if and only if $P(1/f^{k'}) = g \in \mathcal{O}$. That reads $(P - gf^{k'}).(1/f^{k'}) = 0$ and the result follows from Proposition 4.1.

Let us show that $(f^{k'}, \theta + k', (S_{i,j}))$ form an F -involutive basis. We claim that the relations between the symbols $\sigma(f^{k'}) = f^{k'}$, $\sigma(\theta + k') = \sigma(\theta)$, $(\sigma(S_{i,j}))$ are generated by the relations (10), (11), (12) and the following:

$$-\xi_j f^{k'} + f^{k'-1} f'_j \sigma(\theta) - \sum_i f^{k'-1} w_i x_i \sigma(S_{j,i}) = 0 \quad (13)$$

for any $j = 1, \dots, n$. Let $P \in \text{gr}^F(\mathcal{D})$ homogeneous such that

$$Pf^{k'} \in ((\sigma(\theta)), (\sigma(S_{i,j}))).$$

The element P must have degree at least 1, i.e. $P \in (\xi_i)$ and conversely each $P \in (\xi_i)$ is allowed, using (13). The relations between the elements $(\theta + k, (S_{i,j}))$ are generated by (10), (11), (12), which proves our claim. We then conclude to prove that $(f^k, \theta + k, (S_{i,j}))$ form an F -involutive basis by saying that the relations given above can be lifted: we already mentioned it for (10), (11), (12); and (13) is lifted as follows:

$$-\partial_j f^{k'} + f^{k'-1} f'_j (\theta + k') - \sum_i f^{k'-1} w_i x_i S_{j,i} = 0.$$

Finally, these computations prove that the presentation is minimal. □

4.2 Another presentation of $\mathcal{O}[1/f]/\mathcal{O}$

We still assume that f is quasi-homogeneous with an isolated singularity at the origin, and we will derive from Proposition 3.2 and our results in [1] another minimal presentation of the module $N = \mathcal{O}[1/f]/\mathcal{O}$. Let $\mathcal{O}_{x,t} = \mathbb{C}\{\mathbf{x}, t\}$. We will apply Proposition 3.2 to the $\mathcal{D}_{x,t}$ -module

$$M = \frac{\mathcal{O}_{x,t}[\frac{1}{f-t}]}{\mathcal{O}_{x,t}}.$$

which is quasi-isomorphic to the algebraic local cohomology $\mathbb{R}^{\bullet+1}\Gamma_{[f-t=0]}(\mathcal{O}_{x,t})$. There is a $\mathcal{D}_{x,t}$ -module structure on $\mathcal{O}[1/f, s]f^s$ such that s acts as $-\partial_t t$ and $\mathcal{D}_{x,t}f^s \simeq M$ by the mapping $f^s \mapsto [1/(f-t)]$. We endow M with the good bifiltration

$$F_{d,k}(M) = F_{d,k}(\mathcal{D}_{x,t})[1/(f-t)].$$

Lemma 4.1. *M satisfies the assumptions in Proposition 3.3:*

1. $t : \text{gr}^F(M) \rightarrow \text{gr}^F(M)$ is injective,
2. For any d, k , $F_{d,k}(M) = F_d(M) \cap V_k(M)$.

Proof. By e.g. [1], Lemma 2.1., we have

$$\mathrm{gr}^F(M) \simeq \frac{\mathrm{gr}^F(\mathcal{D}_{x,t})}{(t-f, (\xi_i + f'_i \tau)_i)},$$

then 1. comes from the fact that $(t, t-f, (\xi_i + f'_i \tau)_i)$ form a regular sequence. Finally, [1], Proposition 3.2, Proposition 2.6 and Lemma 1.1 give 2. \square

Lemma 4.2. *There is an isomorphism $M/tM \simeq N$ defined as follows: if $g(x, t) \in \mathcal{O}_{x,t}$, then $[g(x, t)/(f-t)^r]$ is mapped to $[g(x, 0)/f^r]$.*

The proof is straightforward. Let us note, for example, that the image of $\partial_t^i \cdot [1/(f-t)]$ is $(j!) [1/f^{j+1}]$.

The b -function of M as defined by Oaku-Takayama, denoted by $b_M(X)$, and the Bernstein-Sato polynomial $b_f(s)$ of f are very close. By definition, $b_f(s)$ satisfies

$$b_f(s) \mathcal{D}[s] f^s \subset \mathcal{D}[s] f^{s+1}.$$

On the other hand, we have $V_0(M) = \mathcal{D}[s] f^s$ and $V_{-1}(M) = \mathcal{D}[s] f^{s+1}$, thus $b_M(X)$ is defined by

$$b_M(t \partial_t) \mathcal{D}[s] f^s \subset \mathcal{D}[s] f^{s+1}.$$

Identifying s and $-\partial_t t$, we get $b_M(-(s+1)) = b_f(s)$, and finally $k' = k_1 + 1$.

From now on we endow the module N with the following good F -filtration, which takes into account the order of the pole f :

$$F_d(N) = \sum_{1 \leq j \leq k'} F_{d-(j-1)}(\mathcal{D}) \left[\frac{1}{f^j} \right],$$

i.e. $\mathrm{ord}^F(1/f^j) = j-1$ for any $1 \leq j \leq k'$. In particular, the identity $1/f^j = f \cdot 1/f^{j+1}$ for $1 \leq j \leq k'-1$ is not adapted to that filtration.

Proposition 4.3. *The module $N = \mathcal{O}[1/f]/\mathcal{O}$ admits the minimal filtered presentation $\mathcal{L}_1 \xrightarrow{\varphi_1} \mathcal{L}_0 \xrightarrow{\varphi_0} N \rightarrow 0$ where*

$$\mathcal{L}_0 = \bigoplus_{k=0}^{k_1} \mathcal{D} \partial_t^k \text{ with } \mathrm{ord}^F(\partial_t^k) = k,$$

$$\varphi_0(\partial_t^k) = (k!) [1/f^{k+1}],$$

$$\mathcal{L}_1 = (\bigoplus_{k=0}^{k_1} \mathcal{D} \partial_t^k X_1) \oplus (\bigoplus_{i=1}^n \bigoplus_{k=0}^{k_1-1} \mathcal{D} \partial_t^k e_i) \oplus \mathcal{D} X_2$$

$$\text{with } \mathrm{ord}^F(\partial_t^k X_1) = k+1, \mathrm{ord}^F(\partial_t^k e_i) = k+1, \mathrm{ord}^F(X_2) = 0,$$

$$\varphi_1(\partial_t^k X_1) = (k+1+\theta) \partial_t^k,$$

$$\varphi_1(\partial_t^k e_i) = -(f'_i \partial_t^{k+1} + \partial_i \partial_t^k),$$

$$\varphi_1(X_2) = f.$$

Proof. Let $S_1 = \mathbb{C}X_1 \oplus \mathbb{C}X_2$ and $\Lambda^j = \Lambda^j \mathbb{C}^n$ with basis $e_{i_1} \wedge \cdots \wedge e_{i_j}$ with $1 \leq i_1 < \cdots < i_j \leq n$. In the sequel, the tensor products are understood over \mathbb{C} . Let δ_1 (resp. δ_2) denote the Koszul map associated with the sequence f'_1, \dots, f_n (resp. $-\partial_1, \dots, -\partial_n$). For $i = 1, 2$, δ_i is a map $\mathcal{D} \otimes \Lambda^\bullet \rightarrow \mathcal{D} \otimes \Lambda^{\bullet-1}$ and can be extended to a map $\mathcal{D}_{x,t} \otimes \Lambda^\bullet \rightarrow \mathcal{D}_{x,t} \otimes \Lambda^{\bullet-1}$ or to a map $\mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^\bullet \rightarrow \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^{\bullet-1}$ as well.

Our starting point is the beginning of the bifiltered free resolution of M , lifted from the bigraded resolution computed in [1], proof of Theorem 3.2:

$$\mathcal{L}_2 \xrightarrow{\psi_2} \mathcal{L}_1 \xrightarrow{\psi_1} \mathcal{D}_{x,t}[0][0] \xrightarrow{\psi_0} M \rightarrow 0$$

where

$$\psi_0(1) = [1/(f-t)];$$

$$\mathcal{L}_1 = (\mathcal{D}_{x,t} \otimes S_1) \oplus (\mathcal{D}_{x,t} \otimes \Lambda^1) \oplus (\mathcal{D}_{x,t} \otimes \Lambda^2)$$

$$\text{with } \text{ord}^{F,V}(X_1) = (1, 0); \text{ord}^{F,V}(X_2) = (0, 0); \text{ord}^{F,V}(e_i) = (1, 1); \\ \text{ord}^{F,V}(e_i \wedge e_j) = (1, 0);$$

$$\psi_1(X_1) = \partial_t t + \theta; \psi_1(X_2) = f - t; \psi_1(e_i) = -f'_i \partial_t - \partial_i; \psi_1(e_i \wedge e_j) = -S_{i,j};$$

$$\mathcal{L}_2 = \mathcal{D}_{x,t} \cdot 1 \oplus (\mathcal{D}_{x,t} \otimes \Lambda^1) \oplus (\mathcal{D}_{x,t} \otimes \Lambda^2 \otimes S_1) \oplus (\mathcal{D}_{x,t} \otimes \Lambda^3 \otimes S_1)$$

$$\text{with } \text{ord}^{F,V}(1) = (1, 1); \text{ord}^{F,V}(e_i) = (1, 0); \text{ord}^{F,V}(e_i \wedge e_j \otimes X_1) = (1, 1); \\ \text{ord}^{F,V}(e_i \wedge e_j \otimes X_2) = (2, 1); \text{ord}^{F,V}(e_i \wedge e_j \wedge e_k \otimes X_1) = (1, 0); \text{ord}^{F,V}(e_i \wedge e_j \wedge e_k \otimes X_2) = (2, 0);$$

$$\psi_2(1) = X_1 + \partial_t X_2 + \sum w_j x_j e_j; \psi_2(e_i) = f'_i X_1 - \partial_i X_2 + t e_i + \sum_j w_j x_j e_i \wedge e_j;$$

$$\psi_2(e_i \wedge e_j \otimes X_1) = \delta_1(e_i \wedge e_j) + e_i \wedge e_j; \psi_2(e_i \wedge e_j \otimes X_2) = \delta_2(e_i \wedge e_j) - \partial_t e_i \wedge e_j;$$

$$\text{for } I = (i, j, k), \psi_2(e_I \otimes X_l) = \delta_l(e_I).$$

By homogenizing with respect to F , this gives the beginning of a minimal V -adapted free resolution

$$\mathbf{R}\mathcal{L}_2 \xrightarrow{\mathbf{R}\psi_2} \mathbf{R}\mathcal{L}_1 \xrightarrow{\mathbf{R}\psi_1} \mathcal{D}_{x,t}^{(h)}[0][0] \xrightarrow{\mathbf{R}\psi_0} \mathbf{R}M \rightarrow 0$$

with the maps $\mathbf{R}\psi_i$ for $i = 1, 2$ having the same expressions as the maps ψ_i .

Then we apply Proposition 3.2, we obtain the following exact sequence:

$$\frac{V_{k_1}(\mathbf{R}\mathcal{L}_2)}{tV_{k_1+1}(\mathbf{R}\mathcal{L}_2)} \xrightarrow{\overline{\mathbf{R}\psi_2}} \frac{V_{k_1}(\mathbf{R}\mathcal{L}_1)}{tV_{k_1+1}(\mathbf{R}\mathcal{L}_1)} \xrightarrow{\overline{\mathbf{R}\psi_1}} \frac{V_{k_1}(\mathcal{D}_{x,t}^{(h)})}{tV_{k_1+1}(\mathcal{D}_{x,t}^{(h)})} \xrightarrow{\overline{\mathbf{R}\psi_0}} \mathbf{R}N \rightarrow 0.$$

We have

$$\mathcal{F}_0 := \frac{V_{k_1}(\mathcal{D}_{x,t}^{(h)})}{tV_{k_1+1}(\mathcal{D}_{x,t}^{(h)})} \simeq \bigoplus_{k=0}^{k_1} \mathcal{D}^{(h)} \partial_t^k$$

with $\deg(\partial_t^k) = k$;

$$\mathcal{F}_1 := \frac{V_{k_1}(\mathbf{RL}_1)}{tV_{k_1+1}(\mathbf{RL}_1)} \simeq \bigoplus_{k=0}^{k_1} \mathcal{D}^{(h)} \partial_t^k \otimes S_1 \oplus \left(\bigoplus_{k=0}^{k_1-1} \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^1 \right) \oplus \left(\bigoplus_{k=0}^{k_1} \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^2 \right)$$

with $\deg(\partial_t^k \otimes X_1) = k + 1$; $\deg(\partial_t^k \otimes X_2) = k + 1$; $\deg(\partial_t^k \otimes e_i) = k + 1$;
 $\deg(\partial_t^k \otimes e_i \wedge e_j) = k + 1$;

$$\begin{aligned} \mathcal{F}_2 &:= \frac{V_{k_1}(\mathbf{RL}_2)}{tV_{k_1+1}(\mathbf{RL}_2)} \\ &\simeq \left(\bigoplus_{k=0}^{k_1-1} \mathcal{D}^{(h)} \partial_t^k \right) \oplus \left(\bigoplus_{k=0}^{k_1} \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^1 \right) \oplus \left(\bigoplus_{k=0}^{k_1-1} \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^2 \otimes S_1 \right) \\ &\quad \oplus \left(\bigoplus_{k=0}^{k_1} \mathcal{D}^{(h)} \partial_t^k \otimes \Lambda^3 \otimes S_1 \right) \end{aligned}$$

with $\deg(\partial_t^k \cdot 1) = k + 1$; $\deg(\partial_t^k \otimes e_i) = k + 1$; $\deg(\partial_t^k \otimes e_i \wedge e_j \otimes X_1) = k + 1$;
 $\deg(\partial_t^k \otimes e_i \wedge e_j \otimes X_2) = k + 2$; for $I = (i, j, k)$, $\deg(\partial_t^k \otimes e_I \otimes X_1) = k + 1$;
 $\deg(\partial_t^k \otimes e_I \otimes X_2) = k + 2$.

Let us compute the maps $\overline{\mathbf{R}\psi}_i$. We have $\overline{\mathbf{R}\psi}_0(\partial_t^k) = (k!)[1/f^{k+1}]T^k$. In $\mathcal{D}_{x,t}^{(h)}$ the following identity holds:

$$\partial_t^k t = t \partial_t^k + k h \partial_t^{k-1}.$$

Then

$$\begin{aligned} \overline{\mathbf{R}\psi}_1(\partial_t^k \otimes X_1) &= [\partial_t^k(\partial_t + \theta)] \\ &= [t \partial_t^{k+1} t + ((k+1)h + \theta) \partial_t^k] \\ &= ((k+1)h + \theta) \partial_t^k. \end{aligned}$$

In the same way the computation gives:

$$\begin{aligned} \overline{\mathbf{R}\psi}_1(\partial_t^k \otimes X_2) &= f \partial_t^k - k h \partial_t^{k-1} \\ \overline{\mathbf{R}\psi}_1(\partial_t^k \otimes e_i) &= -(f'_i \partial_t^{k+1} + \partial_i \partial_t^k) \\ \overline{\mathbf{R}\psi}_1(\partial_t^k \otimes e_i \wedge e_j) &= -S_{i,j} \partial_t^k \\ \overline{\mathbf{R}\psi}_2(\partial_t^k \cdot 1) &= \partial_t^k \otimes X_1 + \partial_t^{k+1} \otimes X_2 + (\sum_j w_j x_j \partial_t^k \otimes e_j) \\ \overline{\mathbf{R}\psi}_2(\partial_t^k \otimes e_i) &= f'_i \partial_t^k \otimes X_1 - \partial_i \partial_t^k \otimes X_2 + k h \partial_t^{k-1} \otimes e_i + \sum_j w_j x_j \partial_t^k \otimes e_i \wedge e_j \\ \overline{\mathbf{R}\psi}_2(\partial_t^k \otimes e_i \wedge e_j \otimes X_1) &= \delta_1(\partial_t^k \otimes e_i \wedge e_j) + \partial_t^k e_i \wedge e_j \\ \overline{\mathbf{R}\psi}_2(\partial_t^k \otimes e_i \wedge e_j \otimes X_2) &= \delta_2(\partial_t^k \otimes e_i \wedge e_j) - \partial_t^{k+1} e_i \wedge e_j \\ \overline{\mathbf{R}\psi}_2(\partial_t^k \otimes e_I \otimes X_I) &= \delta_I(\partial_t^k \otimes e_I). \end{aligned}$$

Since some of the entries of the matrix which represents $\overline{\mathbf{R}\psi_2}$ are units, we have to minimalize. Our method is the same as the minimalization process described in [6], paragraph 4.4. Let

$$\mathcal{G} = \bigoplus_{k=0}^{k_1-1} \mathcal{D}^{(h)} \partial_t^k \oplus \bigoplus_{i < j} \bigoplus_{k=0}^{k_1-1} \mathcal{D}^{(h)} \partial_t^k \otimes e_i \wedge e_j \otimes X_2 \oplus \bigoplus_{i < j} \mathcal{D}^{(h)} \otimes e_i \wedge e_j \otimes X_1,$$

a free submodule of \mathcal{F}_2 . We have the following commutative diagram:

$$\begin{array}{ccccc} 0 & & 0 & & \\ \downarrow & & \downarrow & & \\ \mathcal{G} & \xrightarrow{\overline{\mathbf{R}\psi_2}} & \overline{\mathbf{R}\psi_2}(\mathcal{G}) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{F}_2 & \xrightarrow{\overline{\mathbf{R}\psi_2}} & \mathcal{F}_1 & \xrightarrow{\overline{\mathbf{R}\psi_1}} & \mathcal{F}_0 \\ \downarrow & & \downarrow & & \downarrow \text{Id} \\ \mathcal{F}_2/\mathcal{G} & \xrightarrow{\phi_2} & \mathcal{F}_1/\overline{\mathbf{R}\psi_2}(\mathcal{G}) & \xrightarrow{\phi_1} & \mathcal{F}_0 \\ \downarrow & & \downarrow & & \downarrow \\ 0 & & 0 & & 0 \end{array}$$

where ϕ_i is the map induced by $\overline{\mathbf{R}\psi_i}$ for $i = 1, 2$. The columns are exact and the first two rows are exact. Then we deduce that the third row is also exact. This row will provide our minimal presentation. The module $\mathcal{F}'_2 := \mathcal{F}_2/\mathcal{G}$ is free with basis $(\partial_t^k \otimes e_i)_{i,k}$, $(\partial_t^k \otimes e_i \wedge e_j \otimes X_1)_{k \geq 1, i, j}$ and $(\partial_t^k \otimes e_I \otimes X_j)_{I=(i_1, i_2, i_3), k, j=1, 2}$. It is easy to show that

$$\mathcal{F}'_1 := \frac{\mathcal{F}_1}{\overline{\mathbf{R}\psi_2}(\mathcal{G})}$$

is free with basis $(\partial_t^k \otimes X_1)_k$, $(\partial_t^k \otimes e_i)_{i,k}$ and X_2 . Note that in \mathcal{F}'_1 , we have

$$\begin{aligned} e_i \wedge e_j &\equiv -\delta_1(e_i \wedge e_j), \\ \partial_t^k \otimes e_i \wedge e_j &\equiv \delta_2(\partial_t^{k-1} \otimes e_i \wedge e_j) \quad \text{for } k \geq 1, \\ \partial_t^k \otimes X_2 &\equiv -(\partial_t^{k-1} \otimes X_1 + \sum_j w_j x_j e_j \partial_t^{k-1}) \quad \text{for } k \geq 1. \end{aligned}$$

Using these identities, we compute the matrix representing ϕ_2 in the basis described above:

For $k \geq 1$,

$$\phi_2(\partial_t^k \otimes e_i) = f'_1 \partial_t^k \otimes X_1 + \partial_i \partial_t^{k-1} \otimes X_1 + (k + w_i) h \partial_t^{k-1} \otimes e_i + \sum_j w_j x_j \partial_j \partial_t^{k-1} \otimes e_i;$$

$$\phi_2(e_i) = f'_i X_1 - \partial_i X_2 - \sum_j w_j x_j \delta_1(e_i \wedge e_j);$$

For $k \geq 1$, $\phi_2(\partial_t^k \otimes e_i \wedge e_j \otimes X_1) = \delta_1(\partial_t^k \otimes e_i \wedge e_j) + \delta_2(\partial_t^{k-1} \otimes e_i \wedge e_j)$.

Let $I = (i_1, i_2, i_3)$.

For $k \geq 1$, $\phi_2(\partial_t^k \otimes e_I \otimes X_1) = \delta_2(\delta_1(\partial_t^{k-1} \otimes e_I))$;

$\phi_2(e_I \otimes X_1) = -\delta_1(\delta_1(e_I)) = 0$;

For $k \geq 1$, $\phi_2(\partial_t^k \otimes e_I \otimes X_2) = 0$;

$\phi_2(e_I \otimes X_2) = -\delta_1(\delta_2(e_I))$.

Thus the matrix representing ϕ_2 does not contain any unity. By dehomogenizing we obtain the announced minimal presentation of N . \square

Acknowledgements

I sincerely thank Toshinori Oaku for having suggested the problem discussed in section 3, and for his comments. I am also grateful to Michel Granger for his advices throughout the elaboration of that paper. The project was funded by the Japan Society for the Promotion of Science, The Institute of Mathematics of the University of Seville IMUS, the University of Seville (Spain) and the Erwin Schrödinger International Institute for Mathematical Physics (Vienna, Austria).

References

- [1] R. Arcadias, Minimal resolutions of geometric \mathcal{D} -modules, J. Pure App. Algebra 214 (2010) 1477–1496.
- [2] R. Arcadias, Multidegree for bifiltered D -modules, J. Pure App. Algebra 216 (2012) 280–295.
- [3] F.J. Calderón Moreno, L. Narváez Macarro, The module $\mathcal{D}f^s$ for locally quasi-homogeneous free divisors, Compositio Math. 134 (2002), 59–74.
- [4] F.J. Castro-Jiménez and L. Narváez-Macarro, Homogenising differential operators, Preprint no 36, Universidad de Sevilla, 1997.
- [5] M. Granger and P. Maisonobe, A basic course on differential modules, in: \mathcal{D} -modules cohérents et holonomes, Travaux en cours, vol. 45, Hermann, 1993.
- [6] M. Granger, T. Oaku, Toshinori Minimal filtered free resolutions for analytic D -modules. J. Pure Appl. Algebra 191 (2004), no. 1-2, 157–180.
- [7] M. Kashiwara, P. Schapira, Sheaves on manifolds, With a chapter in French by Christian Houzel, Corrected reprint of the 1990 original, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 292. Springer-Verlag, Berlin, 1994.

- [8] G. Laumon, Sur la catégorie dérivée des \mathcal{D} -modules filtrés, Algebraic geometry (Tokyo/Kyoto, 1982), 151 – 237, Lecture Notes in Math., 1016, Springer, Berlin, 1983.
- [9] T. Oaku, Algorithms for the b -function and D -modules associated with a polynomial, J. Pure App. Algebra 117-118 (1997) 495–518.
- [10] T. Oaku, N. Takayama, Algorithms for D -modules—restriction, tensor product, localization, and local cohomology groups, J. Pure Appl. Algebra 156 (2001), no. 2-3, 267–308.
- [11] T. Oaku, N. Takayama, Minimal free resolutions of homogenized D -modules, Effective methods in rings of differential operators. J. Symbolic Comput. 32 (2001), no. 6, 575–595.
- [12] M. Saito, Modules de Hodge polarisables, Publ. Res. Inst. Math. Sci. 24, No.6, 849-995 (1988).
- [13] U. Walther, Algorithmic computation of De Rham cohomology of complements of complex affine varieties, J. Symbolic Computation 29 (2000), 795–839.